CONTENTS

| At the Cutting Edge | |
|--|------|
| Cretinism, thyroid hormones and selenium | |
| B. Contempré, J. Vanderpas and J.E. Dumont (Belgium) | C193 |
| Diversity in cellular signaling for nerve growth factor and insulin: variations on a common theme | |
| A.R. Saltiel and S.J. Decker (U.S.A.) | C197 |
| Research Papers | |
| Angiotensin II induces calcium release in a subpopulation of single ovarian (granulosa) cells | |
| J.A. Flores, J.D. Veldhuis and D.A. Leong (U.S.A.) | 1 |
| Mechanisms involved in the interaction of dopamine with angiotensin II on al- dosterone secretion in isolated and cultured rat adrenal glomerulosa cells | |
| N. Gallo-Payet, L. Chouinard, MN. Balestre and G. Guillon (Canada, France) Characterization of a 1,25-dihydroxy-vitamin D ₃ receptor in FRTL-5 cells. Evidence | 11 |
| for an inhibitory effect of 1,25-dihydroxy-vitamin D ₃ on thyrotropin-induced iodide uptake | |
| C. Lamberg-Allardt, E. Valtonen, M. Polojärvi and P. Stewen (Finland) | . 25 |
| Angiotensin-II activation of cAMP and corticosterone production in bovine adreno- cortical cells: effects of nonpeptide angiotensin-II antagonists | |
| W.E. Rainey, E.W. Byrd, R.A. Sinnokrot and B.R. Carr (U.S.A.) | 33 |
| Opposite effects of angiotensin-II and corticotropin on bovine adrenocortical cell steroidogenic responsiveness | |
| R. Ouali, D. Langlois, J.M. Saez and M. Begeot (France) | 43 |
| Progesterone-induced second messengers at the onset of meiotic maturation in the amphibian oocyte: interrelationships between phospholipid N-methylation, calcium and diacylglycerol release, and inositol phospholipid turnover | |
| E.J. Chien, G.A. Morrill and A.B. Kostellow (U.S.A.) | 53 |
| Tumor-promoting phorbol ester acts synergistically with insulin to induce lutropin receptor expression in rat granulosa cells | |
| Ma. Hattori, M. Takahashi and R. Horiuchi (Japan) | 69 |
| Analyses of ovine corpora lutea for tumor necrosis factor mRNA and bioactivity during prostaglandin-induced luteolysis | |
| I. Ji, R.G. Slaughter, J.A. Ellis, T.H. Ji and W.J. Murdoch (U.S.A.) | 77 |
| Initial characterization of the four promoters of the human insulin-like growth factor II gene | |
| M.A. Van Dijk, F.M.A. Van Schaik, H.J. Bootsma, P. Holthuizen and J.S. Sussenbach (Netherlands) | 81 |
| Effect of epidermal and insulin-like growth factors on vectorial secretion of transferrin by rat Sertoli cells in vitro | |
| J.A. Spaliviero and D.J. Handelsman (Australia) | 95 |
| | 310 |

(continued on p. 218)

Cited in Biological Abstracts; Current Contents / Life Sciences; Excerpta Medica (EMBASE); Index Medicus; Informedicus; Pascal M

Contents (continued)

| Complement C3 synthesis, peroxidase activity and eosinophil chemotaxis in the rat | |
|--|-----|
| uterus: effect of estradiol and testosterone | 105 |
| M.C. Leiva, C.R. Lyttle and P.H. Jellinck (U.S.A., Canada) | 105 |
| Acute action of luteinizing hormone on mouse Leydig cells: accumulation of | |
| mitochondrial phosphoproteins and stimulation of testosterone synthesis | |
| L.F. Epstein and N.R. Orme-Johnson (U.S.A.) | 113 |
| Mechanism of action of spironolactone on cortisol production by guinea pig adreno- | |
| cortical cells | |
| K.A. Rourke, J.M. Bergstrom, I.W. Larson and H.D. Colby (U.S.A.) | 127 |
| Sites of expression and induction of glandular kallikrein gene expression in mice | |
| J.D. Penschow, C.C. Drinkwater, J. Haralambidis and J.P. Coghlan (Australia) | 135 |
| Efficient production of biologically active human prolactin in Escherichia coli | |
| Y. Hiraoka, Y. Nomata, K. Matsuo, N. Tsubota, K. Tanabe and T. Fukasawa | |
| (Japan) | 147 |
| Hormone formation in the isolated fragment 1-171 of human thyroglobulin involves | |
| the couple tyrosine 5 and tyrosine 130 | |
| C. Marriq, PJ. Lejeune, N. Venot and L. Vinet (France) | 155 |
| Monensin-resistant LLC-PK ₁ cell mutants are affected in recycling of the adenylate | |
| cyclase-stimulating vasopressin V ₂ -receptor | |
| D.A. Jans, P. Jans, H. Luzius and F. Fahrenholz (F.R.G.) | 165 |
| ATP, bradykinin, TRH and TSH activate the Ca ²⁺ -phosphatidylinositol cascade of | |
| human thyrocytes in primary culture | |
| E. Raspé, E. Laurent, G. Andry and J.E. Dumont (Belgium) | 175 |
| Hormone responsive elements within the upstream sequences of the rabbit whey acidic | |
| protein (WAP) gene direct chloramphenicol acetyl transferase (CAT) reporter gene | |
| expression in transfected rabbit mammary cells | |
| E. Devinoy, R. Maliénou-N'Gassa, D. Thépot, C. Puissant and L.M. Houdebine | |
| (France) | 185 |
| Phenol red: an inhibitor of thyroglobulin iodination in cultured porcine thyroid cells | 103 |
| D. Gruffat, S. Gonzalvez, J. Mauchamp and O. Chabaud (France) | 195 |
| Regulation of juvenile hormone synthesis in wild-type and apterous mutant Drosophila | 175 |
| M. Altaratz, S.W. Applebaum, D.S. Richard, L.I. Gilbert and D. Segal (Israel, | |
| U.S.A.) | 205 |
| O.B.A.j | 200 |
| Calendar of Meetings | 217 |
| Carolina of Lavolings | |

Notes to Authors